

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	101-19
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 17.9 kDa, a single non-glycosylated polypeptide chain containing 153 amino acids.
Quantity:	2µg /10µg /1000µg
AA Sequence:	LRRCLISTDM HHIEESFQEI KRAIQAKDTF PNVTLSTLE TLQIIKPLDV CCVTKNLLAF YVDRVFKDHQ EPNPKILRKI SSIANSFLYM QKTLRQCQEQ RQCHCRQEAT NATRVIHDNY DQLEVHAAAI KSLGELDVFL AWINKNHEVM SSA
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using human IL-20R α and human IL-20R β co-transfected murine BaF3 pro-B cells is less than 1.5 ng/ml, corresponding to a specific activity of > 6.7 × 10 ⁵ IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 1 EU/µg of rHuIL-19 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 100 mM HAC to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C . Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Human Interleukin-19

Human Interleukin-19 (IL-19) is encoded by the IL19 gene, which is located on the chromosome 1. The protein belongs to the IL-10 family that includes IL-10, IL-20, IL-22, IL-24, and IL-26. As a monomer made of seven amphipathic helices, IL-19 has a helical bundle and shares the same cell surface receptor (IL-20R) with IL-20 and IL-24. It may play some important roles in inflammatory responses because it up-regulates IL-6 and TNF-alpha and induces apoptosis.