

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	103-06
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 16.3 kDa, a single non-glycosylated polypeptide chain containing 149 amino acids.
Quantity:	10µg/50µg/1000µg
AA Sequence:	MQKGDQNPQI AAHVISEASS KTTSVLQWAE KGYTMSNNL VTLKNGKQLT VKRQGLYYIY AQVTFCSNRE ASSQAPFIAS LWLKSPGRFE RILLRAANTH SSAKPCGQQS IHLGGVFELQ PGASVFNVT DPSQVSHGTG FTSFGLLKL
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by the dose-dependent stimulation of IL-8 production by human PBMC is less than 5-10 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.0.
Endotoxin:	Less than 1 EU/µg of rHusCD40L as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-0.3 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Human soluble CD40 Ligand

CD40 ligand is a 261 amino acid type II transmembrane glycoprotein belonging to the TNF family and can be cleaved into two chains: the CD40 ligand membrane form and CD40 ligand soluble form. It is primarily expressed on activated CD4⁺ T lymphocytes, and also found in other types of cells, like NK cells, mast cells, basophils and eosinophils. In total CD40 ligand has three binding partners: CD40, α5β1 integrin and αIIbβ3. It mediates a range of activities on B cells including induction of activation-associated surface antigen, entry into cell cycle, isotype switching and Ig secretion and memory generation. CD40-CD40L interaction also plays important roles in monocyte activation and dendritic cell maturation.