

Recombinant Rhesus Macaque Granulocyte-Macrophage Colony Stimulating Factor (rRhGM-CSF)

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	112-03
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 14.4 kDa, a single non-glycosylated polypeptide chain containing 127 amino acids.
Quantity:	2µg/10µg/1000µg
AA Sequence:	APARSPSPGT QPWEHVNAIQ EARRLLNLSR DTAAEMNKTV EVVSEMFDLQ EPSCLQTRLE LYKQGLQGSL TKLKGPLTMM ASHYKQHCPP TPETSCATQI ITFQSFKENL KDFLLVIPFD CWEPVQE
Purity:	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using human TF-1 cells is less than 0.1 ng/ml, corresponding to a specific activity of > 1.0 × 10 ⁷ IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 1 EU/µg of rRhGM-CSF as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Rhesus Macaque Granulocyte-Macrophage Colony Stimulating Factor

Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF) is secreted by a number of different cell types (including activated T cells, B cells, macrophages, mast cells, endothelial cells and fibroblasts) in response to cytokine or immune and inflammatory stimulation. It was initially characterized as a growth factor that can support the in vitro colony formation of granulocyte-macrophage progenitors and has functions of stimulates the growth and differentiation of hematopoietic precursor cells from various lineages. GM-CSF has also been reported to have a functional role on non-hematopoietic cells and can induce human endothelial cells to migrate and proliferate. Additionally, it can stimulate the proliferation of a number of tumor cell lines, including osteogenic sarcoma, carcinoma and adenocarcinoma cell lines. It is reported that GM-CSF has no biological effects across species.