

ChemWhat Recombinant Murine Fibroblast Growth Factor-9
A brand under Watson (rMuFGF-9)

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	124-09
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 23.4 kDa, a single non-glycosylated polypeptide chain containing 207 amino acids.
Quantity:	2µg/10µg/1000µg
AA Sequence:	MPLGEVGSYF GVQDAVPFGN VPVLPVDSPV LLNDHLGQSE AGGLPRGPAV TDL DHLK GIL RRRQLY CRTG FHLEIFPNGT IQGTRKDHSR FGILEFISIA VGLVSIRGVD SGLYLG MNEK GELYGSEKLT QECVFREQFE ENWYNTYSSN LYKHVDTGRR YYVALNKDGT PREGTRTKRH QKFTHFLPRP VDPDKVPELY KDILSQS
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by thymidine uptake assay using FGF-receptors transfected BaF3 cells is less than 0.5 ng/ml, corresponding to a specific activity of > 2.0 × 10 ⁶ IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2µm filtered concentrated solution in 20 mM Tris, 500 mM NaCl, pH 8.5.
Endotoxin:	Less than 1 EU/µg of rMuFGF-9 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C . Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Murine Fibroblast Growth Factor-9

Fibroblast growth factor 9 (FGF-9) encoded by the FGF-9 gene is a member of the fibroblast growth factor (FGF) family. It plays an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. In addition, this protein was isolated as a secreted factor that exhibits a growth-stimulating effect on cultured glial cells and it is a heparin-binding protein. Furthermore, FGF-9 is a monomer and interacts with FGFR1, FGFR2, FGFR3 and FGFR4. Recombinant mouse FGF-9 is synthesized as a 208 a.a. precursor that contains a 3 a.a. signal sequence. Specifically, The mouse FGF-9 shares 99 % a. a. sequence identity with rat FGF-9.