

Recombinant Human Granulocyte Chemotactic Protein 2/CXCL6 (rHuGCP-2/CXCL6)

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	201-06
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 7.9 kDa, a single non-glycosylated polypeptide chain containing 72 amino acids.
Quantity:	5µg/20µg/1000µg
AA Sequence:	VLTELRCTCL RVTLRVNPKT IGKLQVFPAG PQCSKVEVVA SLKNGKQVCL DPEAPFLKKV IQKILDSGNK KN
Purity:	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human neutrophils is in a concentration range of 10-50 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 0.1 EU/µg of rHuGCP-2/CXCL6 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C . Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Human Granulocyte Chemotactic Protein 2/CXCL6

GCP-2 (granulocyte chemotactic protein-2) is a CXC chemokine. Among human CXC chemokines, GCP-2 is most closely related to ENA-78. The structure and sequence of the genes for human GCP-2 and ENA-78 also exhibit close similarity, suggesting the two genes may have originated from a recent gene duplication. LIX (LPS-induced CXC chemokine) was initially cloned as a gene induced by LPS in mouse fibroblasts. The mouse protein designated GCP-2, because of its amino acid sequence similarity (60%) to human GCP-2, is identical to the LIX protein sequence.