

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	202-01
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 10.2 kDa, a single non-glycosylated polypeptide chain containing 92 amino acids.
Quantity:	5µg /20µg /1000µg
AA Sequence:	GSEVSDKRTC VSLTTQRLPV SRIKTYTITE GSLRAVIFIT KRGLKVCADP QATWVRDVVR SMDRKSNTRN NMIQTKPTGT QQSTNTAVTL TG
Purity:	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration of 10-100 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Endotoxin:	Less than 1 EU/µg of rHuLymphotactin/XCL1 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Human Lymphotactin/XCL1

Chemokine (C motif) ligand (XCL1), as known as lymphotactin, is the only known member of the C-chemokine family and signals through the receptor XCR1, formally known as GPR5. The expression of lymphotactin is abundant in some activated T cells such as activated CD8+ T cells and other class I MHC restricted T cells. It is found in high levels in spleen, thymus, intestine and peripheral blood leukocytes, and at lower levels in lung, prostate gland and ovary. XCL1 induces its chemotactic function by binding to a chemokine receptor called XCR1. Recombinant Human XCL1 which is a single non-glycosylated polypeptide chains containing 92 amino acids and it shares approximately 60 % amino acid sequence homology with the murine and rat protein.