

Recombinant Human Monocyte Chemotactic Protein-4/CCL13 (rHuMCP-4/CCL13)

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	204-13
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 8.6 kDa, a single non-glycosylated polypeptide chain containing 75 amino acids.
Quantity:	5µg /20µg /1000µg
AA Sequence:	QPDALNVPST CCFTFSSKKI SLQRLKSYVI TTSRCPQKAV IFRTKLGKEI CADPKEKWVQ NYMKHLGRKA HTLKT
Purity:	> 96 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human monocytes is in a concentration of 10-100 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 130 mM NaCl.
Endotoxin:	Less than 1 EU/µg of rHuMCP-4/CCL13 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20°C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Human Monocyte Chemotactic Protein-4/CCL13

Human CCL13 is belonging to the CC chemokine family and is encoded by the gene CCL13 in humans. CCL13 (MCP-4) shares 56-61 % sequence identity with MCP-1 (CCL2) and MCP-3 (CCL7) and is 60 % identical to Eotaxin (CCL11). CCL13 was a potent chemoattractant for monocytes and eosinophils and stimulated histamine release from basophils. CCL13 can induce a calcium flux in HEK-293 cells transfected with the receptor CCR2B and CCR3. That shows the function receptors of CCL3 are CCR2B and CCR3.