

**Recombinant Human Hemofiltrate CC  
Chemokine-1/CCL14  
(rHuHCC-1/CCL14)**

**ChemWhat Technical Data Sheet (TDS)**

---

<b>Catalog Number:</b>	204-14
<b>Source:</b>	<i>Escherichia coli</i> .
<b>Molecular Weight:</b>	Approximately 8.4 kDa, a single non-glycosylated polypeptide chain containing 72 amino acids.
<b>Quantity:</b>	2µg /10µg /1000µg
<b>AA Sequence:</b>	TESSSRGPYH PSECCFTYTT YKIPRQRIMD YYETNSQCSK PGIVFITKRG HSVCTNPSDK WVQDYIKDMK EN
<b>Purity:</b>	> 96 % by SDS-PAGE and HPLC analyses.
<b>Biological Activity:</b>	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human monocytes is in a concentration of 5.0-20 ng/ml.
<b>Physical Appearance:</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 100 mM NaCl.
<b>Endotoxin:</b>	Less than 1EU/µg of rHuHCC-1/CCL14 as determined by LAL method.
<b>Reconstitution:</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
<b>Shipping:</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage:</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>
<b>Usage:</b>	<b>ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.</b>

---

***Human Hemofiltrate CC Chemokine-1/CCL14***

Human CCL14 is belonging to the CC chemokine family. It has two isoforms, CCL14a (HCC-1) and CCL14b (HCC-3). The sequence of HCC-3 differs from HCC-1 as follow: 27-27 R→ QTGGKPKVVVKIQLKLVG. CCL14 was first isolated from the hemofiltrate of human patients with chronic renal failure. CCL14 promotes chemotaxis of T lymphocytes, monocytes and eosinophils, and inhibits infection of M-tropic human immunodeficiency virus type 1 and is a ligand for CCR1, CCR3 and CCR5. CCL14 is expressed constitutively in several normal tissues: spleen, liver, skeletal and heart muscle, gut, and bone marrow, present at high concentrations in plasma.