

Recombinant Human Macrophage Inflammatory Protein-5, 68a.a./CCL15 (rHuMIP-5, 68a.a./CCL15)

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	204-15T
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 7.4 kDa, a single non-glycosylated polypeptide chain containing 68 amino acids.
Quantity:	5µg /25µg /1000µg
AA Sequence:	SFHFAADCCT SYISQSIPCS LMKSYFETSS ECSKPGVIFL TTKGRQVCAK PSGPGVQDCM KKLKPYSI
Purity:	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 1.0-10 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 1 × PBS, pH 7.2.
Endotoxin:	Less than 0.1 EU/µg of rHuMIP-5, 68a.a./CCL15 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Human Macrophage Inflammatory Protein-5, 68a.a./CCL15

Human CCL15 is belonging to the CC chemokine family and shares 35 % amino acid homology with human HCC1 (CCL14). CCL15 is most abundant in heart, skeletal muscle and adrenal gland, and low expressed in liver, small intestine, colon, and in certain leukocytes and macrophages of the lung. It is chemotactic for neutrophils, monocytes, and lymphocytes and elicits its effects by binding to cell surface chemokine receptors like CCR1 and CCR3. CCL15 has several cleaved chains. All of them are more potent chemoattractants than CCL15.