

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	244-24
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 10.5 kDa, a single non-glycosylated polypeptide chain consisting of an N-terminal Methionine and the mature rat Eotaxin-2.
Quantity:	5µg /20µg /1000µg
AA Sequence:	MPTGSVTIPS SCCVTFISKK IPVNRVISYQ LANGSICPKA GVIFITKKGH KICTDPKLPW VQKHIKNLDA KRNQPSEGA ALGPKFVIQK LRGNSTKV
Purity:	> 96 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biologically active determined by a chemotaxis bioassay using human peripheral blood eosinophils is in a concentration of 50-250 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 2 × PBS, pH 7.4.
Endotoxin:	Less than 1 EU/µg of rRtEotaxin-2/CCL24 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C . Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Rat Eotaxin-2/CCL24

CCL24, also named MPIF-2, Eotaxin-2 and Ckβ6, is a novel CC chemokine recently identified. It is a secreted protein, encoded by CCL24 gene, and produced by activated monocytes and T lymphocytes. CCL24 signals through the CCR3 receptor and has functions of chemotactic activity for resting T-lymphocytes and eosinophils, but none for monocytes and activated lymphocytes. The plasma levels of CCL24 and the aspirin-exacerbated respiratory disease (such as asthma) morbidity rate have positive correlation.