

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	251-04
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 9.5 kDa, a single non-glycosylated polypeptide chain containing 88 amino acids.
Quantity:	2µg/10µg/1000µg
AA Sequence:	ESSFPATFVP LPADSEGGED EDLQCVCLKT TSGINPRHIS SLEVIGAGTH CPSPQLLATK KTGRKICLDQ QRPLYKKILK KLLDGDES
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human neutrophils is in a concentration of 10-100ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, 500 mM NaCl, pH 7.0.
Endotoxin:	Less than 0.1 EU/µg of rBoPF-4/CXCL4 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20°C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	ChemWhat Limited in UK offers this branded product for research, development or further evaluation purposes. NOT FOR HUMAN USE.

Bovine Platelet Factor-4/CXCL4

Platelet Factor-4 (PF-4) is a small cytokine belonging to the CXC chemokine family that is also known as chemokine (C-X-C motif) ligand 4 (CXCL4). This chemokine is released from alpha-granules of activated platelets during platelet aggregation and neutralizes the anticoagulant effect of heparin. It interacts with a splice variant of the chemokine receptor CXCR3 besides being chemotactic for neutrophils, fibroblasts and monocytes. Recombinant bovine CXCL4 contains 88 amino acids. Specifically, Bovine and sheep CXCL4 share about an 85 % identity.