

Recombinant Human Cu/Zn Superoxide Dismutase, His (rHuCu/Zn SOD, His)

ChemWhat Technical Data Sheet (TDS)

Catalog Number:

401-01H

Source:

Escherichia coli.

Molecular Weight:

Approximately 39.9 kDa, a homodimer, non-glycosylated polypeptide chain containing 2 × 189

amino acids with Met, Gly and 10 × His at N-terminus.

Quantity:

20µg/100µg/1000µg

AA Sequence:

MGHHHHHHH HHSSGHIEGR HMTYARAAAR QARALEATKA VCVLKGDGPV

QGIINFEQKE SNGPVKVWGS IKGLTEGLHG FHVHEFGDNT AGCTSAGPHF NPLSRKHGGP KDEERHVGDL GNVTADKDGV ADVSIEDSVI SLSGDHCIIG

RTLVVHEKAD DLGKGGNEES TKTGNAGSRL ACGVIGIAQ

Purity:

> 95 % by SDS-PAGE and HPLC analyses.

Biological Activity:

Fully biologically active when compared to standard. The potency per mg was determined by

Pyrogallic Acid method and was found to be more than 1.0×10^4 IU/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation: Endotoxin: Lyophilized from a 0.2 μm filtered concentrated solution in PBS, pH 7.4. Less than 1 EU/μg of rHuCu/Zn SOD, His as determined by LAL method.

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1~% BSA to a

concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and

Shipping:

stored at \leq -20 °C. Further dilutions should be made in appropriate buffered solutions. The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature

recommended below.

Stability & Storage:

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

12 months from date of receipt, -20 to -70 °C as supplied.

1 month, 2 to 8 °C under sterile conditions after reconstitution.

3 months, -20 to -70 °C under sterile conditions after reconstitution.

Usage:

ChemWhat Limited in UK offers this branded product for research, development or further

evaluation purposes. NOT FOR HUMAN USE.

Human Cu/Zn Superoxide Dismutase

Superoxide dismutase catalyzes the reaction between superoxide anions and hydrogen to yield molecular oxygen and hydrogen peroxide. Cu/Zn superoxide dismutase also named as SOD1, is an enzyme encoded by the SOD1 gene in humans, located on chromosome 21. The SOD1 binds Cu and Zn ions and is one of three SODs responsible for destroying free superoxide radicals in the body. It has been shown to interact with CCS and Bcl-2. The malfunction of SOD1 may increase the risk of illnesses like agerelated muscle mass loss (sarcopenia), early development of cataracts, macular degeneration, thymic involution, hepatocellular carcinoma, shortened lifespan, keratoconus and amyotrophic lateral sclerosis.

Rev. 08/20/2018 V.3

CHEMWHAT LIMITED

https://www.chemwhat.com

Email: contact@chemwhat.com