

Recombinant Human Interferon-gamma GMP (rHuIFN- γ GMP)

ChemWhat Technical Data Sheet (TDS)

Catalog Number:	GMP-106-06
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 16.9 kDa, a single non-glycosylated polypeptide chain containing 144 amino acids.
Quantity:	5 μ g/100 μ g/1mg
AA Sequence:	MQDPYVKEAE NLKKYFNAGH SDVADNGTLF LGILKNWKEE SDRKIMQSQI VSFYFKLFKN FKDDQSIQKS VETIKEDMNV KFFNSNKKKR DDFEKLTNYS VTDLNVQRKA IHELIQVMAE LSPAAGTGKR KRSQMLFRGR RASQ
Purity:	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as measured in anti-viral assays using human HeLa cells infected with encephalomyocarditis (EMC) virus is 0.15-0.80 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 5.0, with 3 % Trehalose.
Endotoxin:	Less than 0.01 EU/ μ g of rHuIFN- γ GMP as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at \leq -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● A minimum of 12 months when stored at \leq -20 °C as supplied. Refer to lot specific COA for the Use by Date.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This GMP product can be for research use or further manufacturing use.
Quality statement:	The manufacture and testing of this product is in compliance with ICH Q7a guidelines.

